

U.S. Wheat and Barley Scab Initiative
FY00 Final Performance Report (approx. May 00 – April 01)
July 30, 2001

Cover Page

PI:	DavidSchisler
Institution:	USDA-ARS
Address:	Nat. Ctr. For Ag Utilization Res. 1815 N. University St. Peoria, IL 61604-3999
Email:	schislda@mail.ncaur.usda.gov
Phone:	309-681-6284
Fax:	605-688-4452
Year:	FY2000 (approx. May 00 – April 01)
Grant Number:	
Grant Title:	Fusarium Head Blight Research
2000 ARS Award Amount:	\$9,756

Project

Program Area	Project Title	Requested Amount
Chemical & Biological Control	Field testing of scab antagonists on durum wheat.	\$88,300.00
	Requested Total	\$88,300.00 ¹

Principal Investigator

Date

¹ Note: The Requested Total and the Award Amount are not equal.

Project 1: Field testing of scab antagonists on durum wheat.

1. What major problem or issue is being resolved and how are you resolving it?

Fusarium head blight (FHB) of wheat and barley is a devastating disease capable of reducing yield by 20-40% and more when warm moist conditions exist during the time of flowering and early grain fill. Mycotoxin contamination of grain usually accompanies disease outbreaks. Control measures available for combating this disease are limited. Greenhouse and field experiments funded, in part, by the FHB NRI have demonstrated that 7 biocontrol agents that we had previously discovered were effective in reducing FHB on winter and spring wheats. Experiments were conducted, in work outside the scope of this present investigation, to improve antagonist efficacy and reliability in controlling FHB via management of fermentation protocols. Prior to our investigations, little work had been conducted on biologically controlling FHB on durum wheats. The objective of the presently funded research was to test biocontrol inoculum that had been improved via management of the liquid culture fermentation environment, on two durum wheat cultivars in field trials conducted in Peoria, Illinois and Langdon, ND. Field tests of six antagonists were conducted on two durum cultivars (Ben and Renville) at Peoria, IL and Langdon, ND. Disease severity, incidence, yield and DON content of grain data were collected. In experiments conducted beyond the stated objectives, biomass of the six microbial antagonists was produced in two fermentation media that differed in carbon to nitrogen ratio (6.5 vs 11) but not total carbon loading, and field tested in Peoria, IL on durum wheat cultivars Ben and Renville.

2. What were the most significant accomplishments?

Biological control of FHB on two different cultivars of durum wheat was demonstrated at Peoria, IL and Langdon, ND. Additionally, antagonist biomass produced in C:N 6.5 medium was approximately 18% more effective in reducing FHB severity than was inoculum produced in the C:N 11 medium when data was pooled across both cultivars in the Peoria experiment. At Peoria on cultivar Renville, all six antagonists reduced percent disease severity, percent disease incidence and increased 100 kernel weight when produced in one or both C:N media compared to the buffer control. On cultivar Ben, 3 out of 6 antagonists significantly decreased percent disease severity and increased 100 kernel weight when produced in one or both C:N media. Low viable cell counts adversely affected results in Langdon, ND where only 2 of the 6 antagonists significantly reduced FHB severity on cultivars Renville and Ben. In summary, results on these studies confirmed the feasibility of reducing FHB on wheat and specifically on durum wheat cultivars. The result that the C:N ratio of the antagonist production medium can significantly impact biocontrol agent efficacy deserves further study as a method of increasing the efficacy of proven biocontrol agents.

FY00 (approx. May 00 – April 01)

FY00 Final Performance Report

PI: David Schisler

Grant:

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

SCHISLER, D.A., KHAN, N.I., and BOEHM, M.J. 2000. Greenhouse and field testing of antagonists of Fusarium head blight on durum wheat. *Phytopathology* 90:S69.

SCHISLER, D.A., KHAN, N.I., BOEHM, M.J., SLININGER, P.J., and BOTHAST, R.J. 1999. USDA-ARS, Ohio State University Cooperative research on Biologically controlling Fusarium head blight:1. Antagonist selection and testing on durum wheat. Proceedings of the 1999 National Fusarium Head Blight Forum, Sioux Falls, SD.