USDA-ARS/ U.S. Wheat and Barley Scab Initiative FY07 Final Performance Report (approx. May 07 – April 08) July 15, 2008

Cover Page

PI:	Erick De Wolf
Institution:	Kansas State University
Address:	Dept. of Plant Pathology
	4607 Throckmorton PSC
	Manhattan, KS 66506
E-mail:	dewolf1@ksu.edu
Phone:	785-532-3968
Fax:	785-532-5692
Fiscal Year:	2007
USDA-ARS Agreement ID:	59-0790-7-072
USDA-ARS Agreement	Prediction Models and Improved Pre-Harvest Estimates of
Title:	Deoxynivalenol.
FY07 ARS Award Amount:	\$ 34,695

USWBSI Individual Project(s)

USWBSI Research	Project Title	ARS Adjusted Award
Area	Project The	Amount
EEDF	Prediction Models and Improved Pre-Harvest Estimates of	\$34,695
	Deoxynivalenol.	
	Total Award Amount	\$ 34,695

Principal Investigator

Date

^{*} CBCC – Chemical, Biological & Cultural Control

EEDF - Etiology, Epidemiology & Disease Forecasting

FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain

GET - Genetic Engineering & Transformation

HGR - Host Genetics Resources

HGG - Host Genetics & Genomics

IIR – Integrated/Interdisciplinary Research

PGG - Pathogen Genetics & Genomics

VDUN - Variety Development & Uniform Nurseries

Project 1: Prediction Models and Improved Pre-Harvest Estimates of Deoxynivalenol.

1. What major problem or issue is being resolved and how are you resolving it?

Wheat producers need a reliable disease prediction system to help evaluate the risk of disease and potential need for fungicide applications. Fungicides are an important part of the integrated management for head scab, but are not needed when the disease risk is low. Making the correct decision to apply or not apply improve the potential profit for growers, and helps ensure the grain has lowest possible levels of DON.

2. List the most important accomplishment and its impact (how is it being used?). Complete all three sections (repeat sections for each major accomplishment):

Accomplishment:

Disease prediction models were deployed in 24 states where head scab epidemics have occurred.

Accuracy of the prediction models has been stable at approximately 75 to 80% depending on the type of wheat and production region. We are now testing the first generation of a new type model that should allow us to address parts of the pathogen biology that are not currently part of the prediction system. These new models incorporate a module to predict DON for the first time.

Impact:

Wheat growers in 24 states have access to prediction models for head scab. Growers use these models to assess the risk of disease and respond with fungicides if needed or avoid pesticide use when the risk of disease is low.

The new modeling approach should be able to improve the accuracy of disease predictions. The new models will also incorporate DON into the predictions. These enhancements will increase the value of the prediction models as decision tools used by growers.

As a result of that accomplishment, what does your particular clientele, the scientific community, and agriculture as a whole have now that they didn't have before?:

Wheat growers in 24 states have access to prediction models the help them make disease management decisions based on real-time weather information.

FY07 (approx. May 07 – April 08) PI: De Wolf, Erick USDA-ARS Agreement #: 59-0790-7-072

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

- 1. Paul, P.A., Lipps, P.E., De Wolf, E.D., Shaner, G., Buechley, G., Adhikari, T., Ali, S., Stein, J., Osborne, L. and Madden, L.V. 2007. A distributed lag analysis of the relationship between Gibberella zeae inoculum density on wheat spikes and weather variables. Phytopathology 97:1608-1624.
- 2. De Wolf. E.D. and Isard, S.A. 2007. Disease cycle approach to plant disease prediction. Annual Review of Phytopathology 45:9.1-9.18.