

**USDA-ARS/  
U.S. Wheat and Barley Scab Initiative  
FY11 Final Performance Report  
July 13, 2012**

**Cover Page**

<b>PI:</b>	Jose Costa
<b>Institution:</b>	University of Maryland
<b>Address:</b>	Dept. of Plant Science and Landscape Architecture 2102 Plant Sciences Building College Park, MD 20742-4452
<b>E-mail:</b>	costaj@umd.edu
<b>Phone:</b>	301-405-1317
<b>Fax:</b>	301-314-9308
<b>Fiscal Year:</b>	FY11
<b>USDA-ARS Agreement ID:</b>	59-0206-0-059
<b>USDA-ARS Agreement Title:</b>	Development of Wheat with Resistance to Scab Adapted to the Mid-Atlantic.
<b>FY11 USDA-ARS Award Amount:</b>	\$ 60,278

**USWBSI Individual Project(s)**

<b>USWBSI Research Category*</b>	<b>Project Title</b>	<b>ARS Award Amount</b>
VDHR-SWW	Development of Wheat with Resistance to Scab Adapted to the Mid-Atlantic.	\$ 60,278
	<b>Total ARS Award Amount</b>	<b>\$ 60,278</b>



Principal Investigator

7/3/2012

Date

---

\* MGMT – FHB Management  
 FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain  
 GDER – Gene Discovery & Engineering Resistance  
 PBG – Pathogen Biology & Genetics  
 BAR-CP – Barley Coordinated Project  
 DUR-CP – Durum Coordinated Project  
 HWW-CP – Hard Winter Wheat Coordinated Project  
 VDHR – Variety Development & Uniform Nurseries – Sub categories are below:  
 SPR – Spring Wheat Region  
 NWW – Northern Soft Winter Wheat Region  
 SWW – Southern Soft Red Winter Wheat Region

**Project 1:** *Development of Wheat with Resistance to Scab Adapted to the Mid-Atlantic.*

**1. What major problem or issue is being resolved relevant to Fusarium head blight (scab) and how are you resolving it?**

The major problem being addressed is the need to develop rapidly and effectively host resistance to scab (Fusarium Head Blight) from US native and exotic sources into adapted soft red winter wheat (SRWW) germplasm.

Crosses were made in the greenhouse with US native and exotic resistance sources. Several advanced MD lines with 3BS, 5A and 2DL resistance QTL were selected in 2011 for further testing. Additionally, screening of other MD (University of Maryland) wheat advanced lines was conducted under field conditions in an inoculated nursery at Salisbury (MD) in 2011. Conditions favorable for disease development were aided with daily misting before and during wheat flowering. The scab inoculum was scabby corn grain spread in the field a month before flowering. Additionally, the Southern wheat scab and Northern Uniform Scab Screening nurseries that include new experimental lines were also screened for resistance at Salisbury (MD) with artificial inoculation and misting. The Mason Dixon and Uniform Southern Soft Red Winter Wheat nurseries were also tested. Data for all nurseries was obtained for scab incidence, scab severity, Fusarium damaged kernels, seed weight, plant height, heading date, and DON levels.

Additionally, the complete set of genotypes in the MD wheat state test were screened for resistance at Salisbury (MD) with artificial inoculation and misting. Data for the wheat state test was obtained for scab incidence, scab severity, Fusarium damaged kernels, seed weight, plant height, heading date, and DON levels. Results were published online at the UMDcrops website (<http://mdcrops.umd.edu>) and were widely available to wheat growers.

A collaborative project with VA Tech and NC State University is being conducted to map US native resistance to scab in the soft winter wheat genotype MD01W233-06-1. Doubled haploids of the cross MD01W233-06-1/SS8641 were tested under field conditions in North Carolina and Virginia, and Salisbury, Maryland. A preliminary map of the doubled haploid population was made with simple sequence repeat markers with data produced by the USDA National Genotyping Center in Raleigh, NC. A 9K SNP map of this population is being developed in collaboration with Dr. Gina Brown Guedira at the USDA Genotyping center.

Additionally, to map US native resistance in Roane and Jamestown derived lines, we continued a collaboration with Dr. Carl Griffey at VA Tech on an association wheat mapping project by evaluating 294 lines under misting and inoculation in Salisbury, MD. These were evaluated for incidence, severity, Fusarium damaged kernels, seed weight, plant height, heading date, and DON levels.

Furthermore, we evaluated in our misted nursery in Salisbury (MD), the Neuse/AGS2000 population in collaboration with Dr. Paul Murphy at NCSU. These lines were evaluated for incidence, severity, *Fusarium* damaged kernels, seed weight, plant height, heading date, and DON levels.

**2. List the most important accomplishment and its impact (i.e. how is it being used) to minimize the threat of Fusarium head blight or to reduce mycotoxins. Complete both sections (repeat sections for each major accomplishment):**

**Accomplishment:** Incorporation of the 3BS, 5A and 2DL quantitative trait loci (QTL) of resistance to scab from Sumai3 into adapted soft red winter wheat germplasm such as McCormick and SS8641 by marker-assisted backcrossing. F6 selections were made in 2010-2011.

**Impact:** the availability of these soft red winter wheat lines with scab resistance will reduce scab damage in years favorable to scab development.

**Accomplishment:** Evaluation of the complete set of genotypes in the MD wheat state test for Fusarium head blight resistance at Salisbury (MD) under misting/inoculation and scab resistance data published online.

**Impact:** the availability of this information regarding the resistance of currently grown wheat varieties will allow farmers to select varieties based on scab resistance.

**Accomplishment:** Evaluation of wheat genotypes in the Mason Dixon wheat state test and Uniform Southern Soft Red Winter Wheat nursery for Fusarium head blight resistance at Salisbury (MD) under misting/inoculation and scab resistance data distributed to other breeders.

**Impact:** the availability of this information regarding the resistance of advanced wheat lines in the Mason Dixon wheat state test and Uniform Southern Soft Red Winter Wheat will allow breeders to have this information on lines not specifically bred for scab resistance.

**Include below a list of all germplasm or cultivars released with full or partial support of the USWBSI. List the release notice or publication. Briefly describe the level of FHB resistance.**

None in FY11.

**Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.**

Conway, B., Gao, J., Wang, Y., Murphy, J.P., Brown-Guedira, G., Griffey, Y C., Dong, Y., and J. Costa. 2011. Mapping Scab Resistance in the Winter Wheat Line MD01W233-06-1. Proceedings of the National Fusarium Head Blight Forum; 2011 Dec 4-6. Milwaukee, WI. Lexington, KY: University of Kentucky. p. 15.

Kang, J., Clark, A., Van Sanford, D., Griffey, C., G. L. Brown-Guedira, Dong, Y., J. P. Murphy and J. Costa. 2011. Exotic Scab Resistance Quantitative Trait Loci (QTL) Effects on Soft Red Winter Wheat. *Crop Science* 51:924-933.

Sneller, C., Guttieri, M. , Paul, P., Costa, J. and R. Jackwood. 2011. Variation for Resistance to Kernel Infection and Toxin Accumulation in Winter Wheat Infected with *Fusarium graminearum*. Proceedings of the National Fusarium Head Blight Forum; 2011 Dec 4-6. Milwaukee, WI. Lexington, KY: University of Kentucky. p. 57.

Wright, E., Griffey, C., Malla, S., Van Sanford, D., Harrison, S., Murphy, J.P., Costa, J., Milus, G., Johnson, J., McKendry, A., D. Schmale III and N. McMaster. 2011. Family Based Mapping of Fusarium Head Blight Resistance in Soft Wheat Cultivars Roane and Jamestown. Proceedings of the National Fusarium Head Blight Forum; 2011 Dec 4-6. Milwaukee, WI. Lexington, KY: University of Kentucky. p. 60.