

2008

**NORTH AMERICAN BARLEY SCAB EVALUATION
NURSERY (NABSEN) REPORT**

December 2008

Stephen Neate* and Patrick Gross

Dept Plant Pathology North Dakota State University

***address all enquiries regarding this report to Stephen Neate, address enclosed**

Collaborating Scientists

<p>Richard D. Horsley Professor & Barley Breeder Department of Plant Sciences North Dakota State University P.O. Box 5051 Fargo, ND 58105-5051, U.S.A. Phone: (701) 231-8142 Fax: (701) 231-8474 Email: Richard.Horsley@ndsu.edu</p>	<p>Blake Cooper Sr. Manager North American Barley Research Busch-Agricultural Resources Inc. (BARI) 3515 E. County Rd. 52 Ft. Collins, CO 80524 Phone: (970) 472-2327 Fax: (970) 472-2334 email: blake.cooper @ anheuser-busch.com</p>
<p>Stephen Neate Associate Professor Department of Plant Pathology North Dakota State University P.O. Box 5012 Fargo ND 58105-5012 Phone: 1 701 231-7078 Fax: 1 701 231-7851 Email: stephen.neate@ndsu.edu</p>	<p>James Tucker Agriculture & Agri-Food Canada Research Centre P.O. Box 1000A, R.R. #3 Brandon, Manitoba R7A 5Y3 Canada Phone: (204)726-7650 Fax: (204)728-3858 Email: JTucker@AGR.GC.CA</p>
<p>Bill Legge Agriculture & Agri-Food Canada Research Centre P.O. Box 1000A, R.R. #3 Brandon, Manitoba R7A 5Y3 Canada Phone: (204)726-7650 Fax: (204)728-3858 Email: blegge@em.agr.ca</p>	<p>Ruth Dill-Macky Associate Professor Department of Plant Pathology University of Minnesota St Paul MN 55108 Phone: 612-625-2227 Fax: 612-625-9728 Email: ruthdm@umn.edu</p>
<p>Kevin P. Smith Assistant Professor Department of Agronomy and Plant Genetics University of Minnesota St. Paul, MN 55108 phone 612-624-1211 fax 612-625-1268 email: smith376@umn.edu</p>	<p>Linnea G. Skoglund Cereal Pathology Manager Busch Agricultural Resources, Inc. 3515 East County Road 52 Fort Collins, CO 80524 Phone: (970) 472-2332 Fax: (970) 472-2334 Email: Linnea.Skoglund@anheuser-busch.com</p>

INTRODUCTION

The 2008 North American Barley Scab Evaluation Nursery (NABSEN) was grown at Fargo, Langdon, Osnabrock and Casselton, ND; St. Paul and Crookston MN, and Brandon, Manitoba. Nurseries were either misted or unmisted (dryland). Dryland nurseries provide conditions similar to those found in commercial fields. Disease in misted fields are more severe than growers would observe in most years and entries with only moderate FHB resistance have high disease levels. Only entries with higher levels of resistance similar to Chevron or CIho 4196 are scored as resistant in the misted nurseries. Dryland nurseries allow discrimination of entries with moderate to low levels of FHB resistance. Each nursery included a set of common checks. The checks were CIho 4196 (resistant two-row check), Chevron (resistant six-row check), Robust and Stander (susceptible six-row checks), MNBrite (moderately resistant six-row check), and Conlon (moderately resistant two-row check). At all locations percent severity of FHB was determined at the soft to middle dough stage by determining the ratio of infected kernels to total kernels on 10-20 spikes per entry, and then multiplying by 100.

Site details are as follows;

FARGO, ND – Stephen Neate and Patrick Gross

- Misted
- Inoculated by grain spawn method
- 3 Replicates
- Disease severity - percentage of infected kernels
- Disease incidence - percentage of infected heads
- DON content (ppm) measured by GC/ECD by P Schwarz, NDSU on a composite sample of 3 replicates
- Day to heading counted from date planted to 50% of heads emerged 50%

LANGDON, ND – Stephen Neate and Patrick Gross

- Misted
- Inoculated by grain spawn method
- 3 Replicates
- Disease severity - percentage of infected kernels
- Disease incidence - percentage of infected heads
- DON content (ppm) measured by GC/ECD by P Schwarz, NDSU on a composite sample of 3 replicates
- Day to heading counted from date planted to 50% of heads emerged 50%

OSNABROCK, ND – Richard Horsley

- Dryland
- Inoculated by grain spawn method
- 3 Replicates
- Disease severity - percentage of infected kernels
- Disease incidence - percentage of infected heads
- DON content (ppm) measured by GC/ECD by P Schwarz, NDSU on a composite sample of 3 replicates

Casselton, ND - Linnea Skoglund

- Dryland
- No data

ST. PAUL, MN– Kevin Smith and Ruth Dill-Macky

- Misted
- No data

CROOKSTON, MN – Kevin Smith and Ruth Dill-Macky

- Separate Misted and Dryland trails (only data for Misted)
- Inoculated by spore spray method
- Disease severity - percentage of infected kernels
- Disease incidence - percentage of infected heads
- DON content (ppm) measured by GC/ECD by P Schwarz, NDSU on a composite sample of 3 replicates

BRANDON, MANITOBA - Bill Legge and James Tucker

- Misted
- 4 replicates RCB design
- Disease severity - percentage of infected kernels
- Disease incidence - percentage of infected heads
- DON content (ppm) measured by ELISA technique at ECORC, Ottawa on a composite sample of 4 replicates

Table 1. Mean FHB severity of entries grown in the 2008 NABSEN Nursery at five locations.....	6-7
Table 2. Mean disease incidence of entries grown in the 2008 NABSEN Nursery at five locations.....	8-9
Table 3. Mean days to heading after planting of entries grown in 2008 NABSEN Nursery at four locations.....	10-11
Table 4. Mean for DON(ppm) entries grown in 2008 NABSEN Nursery at three locations.....	12-13
Table 5. Average means of Heading date. FHB incidence, FHB severity and DON content.....	14-15
Table 6. Correlation among locations for FHB severity in 2008.....	16
Table 7. Correlation among locations for FHB incidence in 2008.....	16
Table 8. Temperature (°F) compared to the 30-year average.....	17
Table 9. Rainfall (in.) compared to the 30-year average.....	17
Table 10. Pedigree and source of breeding lines tested for FHB resistance in 2008.....	18-19

Table 1. Mean FHB severity of entries grown in the 2008 NABSEN Nursery at five locations.

Label	Brandon	Fargo	Langdon	Crookston		All loc	<u>misted</u>	Osnabrock
				misted	Osnabrock		Mean	dryland
ND23497	23.8	8.8	18.2	11.6	1.1	12.7	15.6	1.1
ND23753	9.9	17.3	23.6	13.6	0.7	13.0	16.1	0.7
ND24978	30.9	6.4	17.5	9.6	0.5	13.0	16.1	0.5
ND25908	30.8	13.9	20.5	26.8	5.7	19.5	23.0	5.7
ND25917	29.7	9.0	9.3	12.2	0.7	12.2	15.0	0.7
ND25977	16.5	16.5	13.7	19.2	1.0	13.4	16.5	1.0
ND26086	15.7	16.1	32.6	26.2	0.9	18.3	22.6	0.9
ND26247	38.0	19.6	21.5	25.3	2.9	21.5	26.1	2.9
Pinnacle	19.9	10.6	16.5	11.3	2.3	12.1	14.6	2.3
2ND28167	25.2	8.4	20.3	30.7	1.9	17.3	21.2	1.9
2ND22182	26.5	17.8	23.6	19.7	3.0	18.1	21.9	3.0
2ND22927	26.9	9.5	8.5	23.5	2.0	14.1	17.1	2.0
2ND24238	20.2	7.1	19.1	27.2	3.2	15.4	18.4	3.2
2ND24263	23.0	11.6	23.2	28.5	1.3	17.5	21.6	1.3
2ND24388	18.8	8.9	25.0	19.7	2.3	14.9	18.1	2.3
2ND24393	22.5	9.8	13.5	16.3	3.8	13.2	15.5	3.8
M128	23.1	11.7	18.4	13.6	1.0	13.5	16.7	1.0
M129	13.2	8.7	16.8	11.2	2.0	10.4	12.5	2.0
M130	22.6	17.9	25.8	11.6	1.1	15.8	19.5	1.1
M132	18.8	16.8	15.3	12.2	1.2	12.8	15.8	1.2
M134 (FEG118-69)	15.2	16.4	15.9	10.2	0.7	11.7	14.4	0.7
M135 (FEG141-20)	17.4	17.7	18.4	11.2	1.8	13.3	16.2	1.8
M136 (FEG147-37)	24.1	10.6	17.3	12.4	2.5	13.4	16.1	2.5
M137 (FEG150-42)	24.0	19.1	25.6	14.1	1.3	16.8	20.7	1.3
6B05-0922	17.6	18.4	22.1	16.8	0.7	15.1	18.7	0.7
6B06-1133	24.4	19.0	26.8	26.2	3.7	20.0	24.1	3.7
6B06-1338	31.6	18.5	25.0	24.0	2.5	20.3	24.8	2.5
6B06-1392	33.7	20.8	20.3	33.1	2.2	22.0	27.0	2.2
6B06-1393	30.1	18.4	19.0	22.3	2.9	18.5	22.5	2.9

Table 1. cont. Mean FHB severity of entries grown in the 2008 NABSEN Nursery at five locations.

Label	Brandon	Fargo	Langdon	<u>Crookston</u>		All loc	<u>misted</u>	<u>Osnabrock</u>
				misted	Osnabrock		Mean	dryland
6B06-1394	22.1	18.3	17.4	16.0	0.9	14.9	18.4	0.9
6B06-1409	24.2	23.4	17.3	21.3	1.3	17.5	21.6	1.3
SB050696	25.5	10.6	6.4	17.3	3.6	12.7	15.0	3.6
SB050716	17.8	10.0	9.4	19.7	1.9	11.8	14.2	1.9
SM060100	17.8	2.7	9.3	11.5	0.7	8.4	10.3	0.7
BM0270D-152	11.2	1.5	9.5	13.0	1.1	7.3	8.8	1.1
BM0331D-357	17.2	2.5	6.5	13.5	0.6	8.1	9.9	0.6
BM0331D-373	11.4	3.2	13.3	18.2	0.3	9.3	11.5	0.3
BM0207D-86	17.3	3.8	14.8	9.8	1.6	9.5	11.4	1.6
BM0264D-60	16.1	6.8	8.0	25.3	1.7	11.6	14.1	1.7
Seebe	17.6	3.4	6.1	11.2	0.6	7.7	9.5	0.6
H94051001	17.1	2.4	1.4	8.0	1.0	6.0	7.2	1.0
H96035002	17.1	2.6	4.3	7.5	1.9	6.7	7.9	1.9
H98011002	17.9	11.9	15.7	21.8	2.2	13.9	16.8	2.2
MnBrite	20.8	12.4	8.5	22.4	1.7	13.1	16.0	1.7
Conlon	12.6	11.4	25.7	21.2	1.4	14.4	17.7	1.4
Robust	27.0	20.8	20.5	28.7	2.1	19.8	24.2	2.1
CIHO 4196	16.9	1.5	1.6	6.2	1.0	5.4	6.5	1.0
Chevron	5.6	1.4	5.3	9.0	0.1	4.3	5.3	0.1
Stander	40.7	23.1	18.0	26.3	2.4	22.1	27.0	2.4

Table 2. Mean disease incidence of entries grown in the 2008 NABSEN Nursery at five locations.

Label	Brandon	Fargo	Langdon	Crookston		All loc	<u>misted</u>	Osnabrock
				misted	Osnabrock		Mean	dryland
ND23497	100.0	86.7	100.0	100.0	36.7	84.7	96.7	36.7
ND23753	97.5	93.3	100.0	100.0	23.3	82.8	97.7	23.3
ND24978	100.0	83.3	100.0	100.0	23.3	81.3	95.8	23.3
ND25908	100.0	96.7	100.0	100.0	63.3	92.0	99.2	63.3
ND25917	100.0	93.3	96.7	100.0	30.0	84.0	97.5	30.0
ND25977	97.5	100.0	100.0	100.0	40.0	87.5	99.4	40.0
ND26086	100.0	100.0	100.0	100.0	30.0	86.0	100.0	30.0
ND26247	100.0	100.0	93.3	100.0	66.7	92.0	98.3	66.7
Pinnacle	87.5	93.3	80.0	80.0	46.7	77.5	85.2	46.7
2ND28167	100.0	90.0	86.7	100.0	40.0	83.3	94.2	40.0
2ND22182	97.5	96.7	96.7	100.0	50.0	88.2	97.7	50.0
2ND22927	97.5	90.0	60.0	100.0	36.7	76.8	86.9	36.7
2ND24238	100.0	83.3	90.0	100.0	60.0	86.7	93.3	60.0
2ND24263	100.0	86.7	86.7	100.0	23.3	79.3	93.3	23.3
2ND24388	92.5	96.7	93.3	100.0	40.0	84.5	95.6	40.0
2ND24393	97.5	93.3	86.7	93.3	53.3	84.8	92.7	53.3
M128	97.5	96.7	96.7	100.0	36.7	85.5	97.7	36.7
M129	95.0	93.3	96.7	96.7	63.3	89.0	95.4	63.3
M130	100.0	93.3	100.0	100.0	43.3	87.3	98.3	43.3
M132	100.0	100.0	96.7	100.0	50.0	89.3	99.2	50.0
M134 (FEG118-69)	100.0	100.0	100.0	93.3	26.7	84.0	98.3	26.7
M135 (FEG141-20)	97.5	100.0	100.0	96.7	63.3	91.5	98.5	63.3
M136 (FEG147-37)	100.0	83.3	96.7	100.0	50.0	86.0	95.0	50.0
M137 (FEG150-42)	97.5	100.0	96.7	100.0	36.7	86.2	98.5	36.7
6B05-0922	97.5	96.7	100.0	100.0	30.0	84.8	98.5	30.0
6B06-1133	100.0	100.0	100.0	100.0	70.0	94.0	100.0	70.0
6B06-1338	100.0	100.0	100.0	100.0	63.3	92.7	100.0	63.3
6B06-1392	100.0	100.0	96.7	100.0	43.3	88.0	99.2	43.3
6B06-1393	100.0	100.0	100.0	100.0	60.0	92.0	100.0	60.0

Table 2. cont. Mean disease incidence of entries grown in the 2008 NABSEN Nursery at five locations.

Label	Brandon	Fargo	Langdon	<u>Crookston</u>		All loc	<u>misted</u>	<u>Osnabrock</u>
				misted	Osnabrock		Mean	dryland
6B06-1394	100.0	100.0	96.7	100.0	23.3	84.0	99.2	23.3
6B06-1409	100.0	100.0	93.3	100.0	50.0	88.7	98.3	50.0
SB050696	100.0	96.7	70.0	96.7	50.0	82.7	90.8	50.0
SB050716	97.5	90.0	76.7	100.0	40.0	80.8	91.0	40.0
SM060100	100.0	53.3	66.7	96.7	16.7	66.7	79.2	16.7
BM0270D-152	85.0	43.3	86.7	100.0	26.7	68.3	78.8	26.7
BM0331D-357	97.5	60.0	53.3	100.0	13.3	64.8	77.7	13.3
BM0331D-373	87.5	63.3	93.3	90.0	6.7	68.2	83.5	6.7
BM0207D-86	97.5	56.7	93.3	100.0	33.3	76.2	86.9	33.3
BM0264D-60	95.0	80.0	83.3	100.0	30.0	77.7	89.6	30.0
Seebe	100.0	70.0	70.0	100.0	16.7	71.3	85.0	16.7
H94051001	100.0	50.0	26.7	96.7	23.3	59.3	68.3	23.3
H96035002	97.5	60.0	60.0	83.3	40.0	68.2	75.2	40.0
H98011002	100.0	90.0	83.3	100.0	46.7	84.0	93.3	46.7
MnBrite	100.0	93.3	90.0	100.0	60.0	88.7	95.8	60.0
Conlon	87.5	100.0	100.0	100.0	26.7	82.8	96.9	26.7
Robust	100.0	96.7	100.0	100.0	56.7	90.7	99.2	56.7
CIHO 4196	87.5	50.0	20.0	76.7	23.3	51.5	58.5	23.3
Chevron	72.5	56.7	63.3	100.0	6.7	59.8	73.1	6.7
Stander	100.0	100.0	100.0	100.0	53.3	90.7	100.0	53.3

Table 3. Mean days to heading after planting of entries grown in 2008 NABSEN Nursery at four locations.

Name	Brandon	Fargo	Langdon	Crookston	
				misted	Mean
ND23497	51.3	61	52	58	55
ND23753	51.5	58	51	58	55
ND24978	55.5	63	56	63	59
ND25908	53.0	61	52	59	56
ND25917	53.0	60	52	60	56
ND25977	52.0	61	52	60	56
ND26086	51.0	59	51	57	54
ND26247	52.0	59	52	58	55
Pinnacle	56.5	61	53	60	58
2ND28167	53.8	61	51	55	55
2ND22182	51.3	58	50	59	54
2ND22927	53.5	59	51	58	55
2ND24238	53.0	61	49	57	55
2ND24263	52.5	58	49	56	54
2ND24388	52.5	57	48	56	53
2ND24393	51.8	57	47	57	53
M128	53.0	60	52	61	57
M129	51.5	58	51	57	54
M130	53.0	60	52	59	56
M132	53.0	59	52	59	56
M134 (FEG118-69)	52.3	59	52	60	56
M135 (FEG141-20)	51.8	60	52	59	56
M136 (FEG147-37)	53.0	61	51	60	56
M137 (FEG150-42)	51.8	59	51	58	55
6B05-0922	53.0	62	52	58	56
6B06-1133	50.3	59	52	58	55
6B06-1338	53.0	62	51	58	56
6B06-1392	52.8	59	54	61	57
6B06-1393	52.5	61	55	60	57

Table 3. cont. Mean days to heading after planting of entries grown in 2008 NABSEN Nursery at four locations.

Name	Brandon	Fargo	Langdon	Crookston		Mean
				misted		
6B06-1394	52.8	61	54	61		57
6B06-1395	50.0	60	53	58		55
6B06-1409	53.0	61	52	60		57
SB050696	52.5	59	49	58		55
SB050716	54.5	60	50	59		56
SM060100	56.3	62	54	63		59
BM0270D-152	57.5	64	55	62		60
BM0331D-357	56.0	63	54	62		59
BM0331D-373	55.3	63	54	63		59
BM0207D-86	54.3	62	55	63		58
BM0264D-60	56.3	63	52	62		58
Seebe	58.3	65	55	65		61
H94051001	55.5	63	54	62		59
H96035002	57.5	65	55	64		60
H98011002	54.8	60	53	59		57
MnBrite	53.5	62	53	58		57
Conlon	50.5	57	50	55		53
Robust	52.5	63	52	58		56
CIHO 4196	59.0	65	53	63		60
Chevron	58.5	65	54	63		60
Stander	53.0	62	52	59		57

Table 4. Mean for DON (ppm) entries grown in 2008 NABSEN Nursery at four locations.

Name	Fargo	Langdon	Brandon	Osnabrock	all loc	misted	dryland
ND23497	5.3	40.4	17.0	2.0	16.2	20.9	2.0
ND23753	5.9	31.6	9.7	2.8	12.5	15.7	2.8
ND24978	8.0	64.6	29.3	4.1	26.5	34.0	4.1
ND25908	8.4	40.0	27.9	3.8	20.0	25.4	3.8
ND25917	5.1	35.2	18.9	3.6	15.7	19.7	3.6
ND25977	6.4	33.1	21.0	2.4	15.7	20.1	2.4
ND26086	7.5	57.2	20.4	5.2	22.6	28.4	5.2
ND26247	8.5	49.7	30.7	5.7	23.6	29.6	5.7
Pinnacle	5.3	33.5	20.8	3.0	15.7	19.9	3.0
2ND28167	4.8	30.4	18.4	2.3	14.0	17.9	2.3
2ND22182	8.5	43.4	17.2	2.2	17.8	23.0	2.2
2ND22927	5.3	26.6	13.4	2.9	12.0	15.1	2.9
2ND24238	4.7	21.0	8.6	2.7	9.3	11.4	2.7
2ND24263	5.4	34.4	13.7	2.6	14.0	17.8	2.6
2ND24388	7.4	30.2	12.6	2.2	13.1	16.7	2.2
2ND24393	6.5	26.8	16.3	1.8	12.9	16.5	1.8
M128	6.7	28.6	22.7	2.8	15.2	19.3	2.8
M129	6.8	41.9	16.4	2.7	16.9	21.7	2.7
M130	8.3	41.0	23.0	3.5	18.9	24.1	3.5
M132	6.6	51.8	15.6	2.6	19.2	24.7	2.6
M134 (FEG118-69)	7.8	35.7	14.8	2.4	15.2	19.4	2.4
M135 (FEG141-20)	6.1	44.1	14.1	2.5	16.7	21.4	2.5
M136 (FEG147-37)	7.7	42.9	28.6	3.6	20.7	26.4	3.6
M137 (FEG150-42)	8.8	49.5	20.9	1.6	20.2	26.4	1.6
6B05-0922	7.0	49.8	22.9	2.8	20.6	26.5	2.8
6B06-1133	9.8	49.5	16.5	2.6	19.6	25.2	2.6
6B06-1338	9.2	37.9	25.0	2.6	18.7	24.0	2.6
6B06-1392	9.5	59.4	27.1	3.5	24.9	32.0	3.5
6B06-1393	7.6	70.0	21.3	5.9	26.2	33.0	5.9

Table 4. cont. Mean for DON (ppm) entries grown in 2008 NABSEN Nursery at four locations.

Name	Fargo	Langdon	Brandon	Osnabrock	all loc	misted	dryland
6B06-1394	10.7	57.7	30.7	3.0	25.5	33.0	3.0
6B06-1395	10.9	59.2	24.4	4.3	24.7	31.5	4.3
6B06-1409	11.6	49.9	25.1	2.6	22.3	28.9	2.6
SB050696	5.7	45.6	15.8	3.5	17.7	22.4	3.5
SB050716	4.1	22.9	13.8	2.6	10.9	13.6	2.6
SM060100	4.9	23.3	23.1	2.4	13.4	17.1	2.4
BM0270D-152	4.4	26.8	15.2	0.8	11.8	15.5	0.8
BM0331D-357	3.3	38.4	17.7	2.5	15.5	19.8	2.5
BM0331D-373	2.6	27.6	13.8	1.2	11.3	14.7	1.2
BM0207D-86	3.4	35.6	19.0	1.5	14.9	19.3	1.5
BM0264D-60	4.6	30.2	13.1	1.0	12.3	16.0	1.0
Seebe	4.6	16.7	15.9	1.7	9.7	12.4	1.7
H94051001	3.2	30.6	13.7	2.0	12.4	15.8	2.0
H96035002	3.8	30.9	20.4	2.6	14.4	18.4	2.6
H98011002	3.9	34.0	18.4	1.6	14.5	18.8	1.6
MnBrite	7.8	40.3	21.7	3.7	18.4	23.3	3.7
Conlon	3.8	22.7	10.3	2.3	9.8	12.2	2.3
Robust	10.5	47.9	23.0	2.7	21.0	27.1	2.7
CIHO 4196	3.6	26.6	26.3	2.7	14.8	18.9	2.7
Chevron	1.9	39.1	13.1	1.1	13.8	18.0	1.1
Stander	15.2	51.2	36.0	3.4	26.5	34.1	3.4

Table 5. Average means of Heading date, FHB Incidence, FHB severity and DON content.

Label	Days to ¹	FHB Incidence ²		FHB SEVERITY ³		DON ppm ⁴	
	head	misted	dryland	misted	dryland	misted	dryland
ND23497	55	96.7	36.7	15.6	1.1	20.9	2.0
ND23753	55	97.7	23.3	16.1	0.7	15.7	2.8
ND24978	59	95.8	23.3	16.1	0.5	34.0	4.1
ND25908	56	99.2	63.3	23.0	5.7	25.4	3.8
ND25917	56	97.5	30.0	15.0	0.7	19.7	3.6
ND25977	56	99.4	40.0	16.5	1.0	20.1	2.4
ND26086	54	100.0	30.0	22.6	0.9	28.4	5.2
ND26247	55	98.3	66.7	26.1	2.9	29.6	5.7
Pinnacle	58	85.2	46.7	14.6	2.3	19.9	3.0
2ND28167	55	94.2	40.0	21.2	1.9	17.9	2.3
2ND22182	54	97.7	50.0	21.9	3.0	23.0	2.2
2ND22927	55	86.9	36.7	17.1	2.0	15.1	2.9
2ND24238	55	93.3	60.0	18.4	3.2	11.4	2.7
2ND24263	54	93.3	23.3	21.6	1.3	17.8	2.6
2ND24388	53	95.6	40.0	18.1	2.3	16.7	2.2
2ND24393	53	92.7	53.3	15.5	3.8	16.5	1.8
M128	57	97.7	36.7	16.7	1.0	19.3	2.8
M129	54	95.4	63.3	12.5	2.0	21.7	2.7
M130	56	98.3	43.3	19.5	1.1	24.1	3.5
M132	56	99.2	50.0	15.8	1.2	24.7	2.6
M134 (FEG118-69)	56	98.3	26.7	14.4	0.7	19.4	2.4
M135 (FEG141-20)	56	98.5	63.3	16.2	1.8	21.4	2.5
M136 (FEG147-37)	56	95.0	50.0	16.1	2.5	26.4	3.6
M137 (FEG150-42)	55	98.5	36.7	20.7	1.3	26.4	1.6
6B05-0922	56	98.5	30.0	18.7	0.7	26.5	2.8
6B06-1133	55	100.0	70.0	24.1	3.7	25.2	2.6
6B06-1338	56	100.0	63.3	24.8	2.5	24.0	2.6
6B06-1392	57	99.2	43.3	27.0	2.2	32.0	3.5
6B06-1393	57	100.0	60.0	22.5	2.9	33.0	5.9

Table 5. cont. Average means of Heading date, FHB Incidence, FHB severity and DON content.

Label	<u>Days to</u> ¹	<u>FHB Incidence</u> ²		<u>FHB SEVERITY</u> ³		<u>DON ppm</u> ⁴	
	head	misted	dryland	misted	dryland	misted	dryland
6B06-1394	57	99.2	23.3	18.4	0.9	33.0	3.0
6B06-1395	55	100.0	70.0	18.2	2.9	31.5	4.3
6B06-1409	57	98.3	50.0	21.6	1.3	28.9	2.6
SB050696	55	90.8	50.0	15.0	3.6	22.4	3.5
SB050716	56	91.0	40.0	14.2	1.9	13.6	2.6
SM060100	59	79.2	16.7	10.3	0.7	17.1	2.4
BM0270D-152	60	78.8	26.7	8.8	1.1	15.5	0.8
BM0331D-357	59	77.7	13.3	9.9	0.6	19.8	2.5
BM0331D-373	59	83.5	6.7	11.5	0.3	14.7	1.2
BM0207D-86	58	86.9	33.3	11.4	1.6	19.3	1.5
BM0264D-60	58	89.6	30.0	14.1	1.7	16.0	1.0
Seebe	61	85.0	16.7	9.5	0.6	12.4	1.7
H94051001	59	68.3	23.3	7.2	1.0	15.8	2.0
H96035002	60	75.2	40.0	7.9	1.9	18.4	2.6
H98011002	57	93.3	46.7	16.8	2.2	18.8	1.6
MnBrite	57	95.8	60.0	16.0	1.7	23.3	3.7
Conlon	53	96.9	26.7	17.7	1.4	12.2	2.3
Robust	56	99.2	56.7	24.2	2.1	27.1	2.7
CIHO 4196	60	58.5	23.3	6.5	1.0	18.9	2.7
Chevron	60	73.1	6.7	5.3	0.1	18.0	1.1
Stander	57	100.0	53.3	27.0	2.4	34.1	3.4

¹ Date from planting to 50% of heads 50% emerged at four locations.

² FHB incidence means at five locations.

³ FHB severity means at five locations.

⁴ DON content means at four locations.

Table 6. Correlation among locations for FHB severity in 2008					
Location	Brandon	Fargo	Langdon	Crookston- misted	Osnabrock
Brandon	1	0.45	0.29	0.35*	0.42*
Fargo	0.45*	1	0.71**	0.35*	0.60**
Langdon	0.29	0.71**	1	0.51**	0.41*
Crookston- Misted	0.35*	0.35*	0.51**	1	0.13
Osnabrock	0.42*	0.60**	0.41*	0.13	1

*,** r-values significantly different from 0.0 at P<0.05 and P<0.01, respectively

Table 7. Correlation among locations for FHB incidence in 2008					
Location	Brandon	Fargo	Langdon	Crookston- misted	Osnabrock
Brandon	1	0.50**	0.25	0.45**	0.46**
Fargo	0.50**	1	0.68**	0.49**	0.29*
Langdon	0.25	0.68**	1	0.48**	0.25
Crookston- Misted	0.45**	0.49**	0.48**	1	0.49**
Osnabrock	0.46**	0.29*	0.25	0.49**	1

*,** r-values significantly different from 0.0 at P<0.05 and P<0.01, respectively

Table 8. Temperature (°F) compared to the 30-year average.

Location	May	June	July	August
Fargo	-3.4	-2.2	0.40	1.00
Langdon	-4.6	-2.5	-0.40	1.80
Crookston, MN	-5.7	-3.3	-1.90	-0.20
Brandon, Manitoba	NA	-2.3	-1.98	1.98

Table 9. Rainfall (in.) compared to the 30-year average.

Location	May	June	July	August
Fargo	-0.51	2.88	-1.23	2.02
Langdon	-1.67	0.96	-0.89	-0.08
Crookston, MN	-1.55	0.57	-0.97	1.6
Brandon, Manitoba	NA	2.45	1.59	-1.27

Table 10. Pedigree and source of breeding lines tested for FHB resistance in 2008.

Label	Pedigree	Contributor	Row type
ND23497	Drummond/ND20414	North Dakota State Univ.	6 row
ND23753	ND18546/ND20407	North Dakota State Univ.	6 row
ND24978	Stellar-ND/6B99-6774	North Dakota State Univ.	6 row
ND25908	ND19655/ND20542	North Dakota State Univ.	6 row
ND25917	ND19655/ND20542	North Dakota State Univ.	6 row
ND25977	ND20303/ND20473	North Dakota State Univ.	6 row
ND26086	ND20477/ND20666	North Dakota State Univ.	6 row
ND26247	ND10655/FEG65-2	North Dakota State Univ.	6 row
Pinnacle	ND18172/ND19130	North Dakota State Univ.	2 row
2ND28167	ND18172/ND19130	North Dakota State Univ.	2 row
2ND22182	ND18413/ND19134//ND19164	North Dakota State Univ.	2 row
2ND22927	ND19119-1/ND19931	North Dakota State Univ.	2 row
2ND24238	ND19854//ND20028/ND19119-1	North Dakota State Univ.	2 row
2ND24263	ND19869/3/ND18998//ND16092/ND17268	North Dakota State Univ.	2 row
2ND24388	ND17274/ND19119//ND19854	North Dakota State Univ.	2 row
2ND24393	ND17274/ND19119//ND19854	North Dakota State Univ.	2 row
M128	FEG26-50 / FEG18-27	University of Minnesota	6 row
M129	FEG59-09 / M110	University of Minnesota	6 row
M130	FEG26-50 / FEG18-27	University of Minnesota	6 row
M132	M96-203 / FEG55-14	University of Minnesota	6 row
M134 (FEG118-69)	FEG69-38 / M114	University of Minnesota	6 row
M135 (FEG141-20)	FEG97-44 / M118	University of Minnesota	6 row
M136 (FEG147-37)	FEG90-31 / M118	University of Minnesota	6 row
M137 (FEG150-42)	ND20493 / M109	University of Minnesota	6 row
6B05-0922	6B99-6557/6B00-1328	BAR - LLC	6 row
6B06-1133	6B98-9022/6B01-2535	BAR - LLC	6 row
6B06-1338	6B99-6517/C02-6922	BAR - LLC	6 row
6B06-1392	6B00-0906/6B99-6557	BAR - LLC	6 row

Table 10. cont. Pedigree and source of breeding lines tested for FHB resistance in 2008

Label	Pedigree	Contributor	Row type
6B06-1394	6B00-0906/6B99-6557	BAR - LLC	6 row
6B06-1395	6B00-0906/6B99-6557	BAR - LLC	6 row
6B06-1409	6B00-1106/6B00-1335	BAR - LLC	6 row
SB050696	SH00749/TR01364	CDC, University of Saskatchewan	2-row
SB050716	SH00749/TR01364	CDC, University of Saskatchewan	2-row
SM060100	SM02353/SM02687	CDC, University of Saskatchewan	2-row
BM0270D-152	TR04282/Newdale	AAFC-Brandon	2-row
BM0331D-357	Conlon/TR03273	AAFC-Brandon	2-row
BM0331D-373	Conlon/TR03273	AAFC-Brandon	2-row
BM0207D-86	TR361/Newdale	AAFC-Brandon	2-row
BM0264D-60	BM9671-23/Newdale	AAFC-Brandon	2-row
Seebe	Masurka//Muller/Heydla	FCDC, Alberta Agriculture and Food	2-row
H94051001	H93126/SEEBE	FCDC, Alberta Agriculture and Food	2-row
H96035002	I94304/TR251	FCDC, Alberta Agriculture and Food	2-row
H98011002	H95009/H97047	FCDC, Alberta Agriculture and Food	2-row
MnBrite	Check	M90-89/M69	6-row
Conlon	Check	BOWMAN*2/DWS1008/ND10232	2-row
Robust	Check	MOREX/MANKER	6-row
CIHO 4196	Check	UNKNOWN	2-row
Chevron	Check	UNKNOWN ROBUST*2/3/CREE/BONANZA//MANKER/4/ROBUST/ BUMBER	6-row
Stander	Check		6-row